



ECS May Demo Overview

Joe Senftle

jsenftle@eos.hitc.com

22 May 1997

Release B Key Milestones



Historical



Investigators Working Group Meeting (Feb. 25-27, 1997)


“Concern about development progress”

Contingency Planning (March 5, 1997)

“Need to demonstrate mission critical functionality ASAP”

SWAMP (April 2-4, 1997)

“May test of functionality”



What It Is

Snapshot of Mission Critical functionality development

- Ingest, Archive, Production, Distribution

Integrates two instrument PGEs

- ASTER Brightness Temperature PGE processes ASTER L1BT input data to produce L2 brightness temperature
- MOPITT CalRads PGE processes L0 input data to produce L1 radiance



DSWG Functions in Demo

Ingest all instrument Level 0 or equivalent source data (ASTER Levels 1A and 1B, Landsat 7 Level 0R, and ancillary data)

- Demo ingests MOPITT Level 0 and ASTER Level 1

Catalog data such that they can be located

- MOPITT and ASTER source data and products are cataloged, searched and retrieved

Run PGEs to support pre- and post-launch testing

- MOPITT Level 1 and ASTER Level 2 PGEs executed in Demo



Support data retrieval by Instrument Team (IT) users

- MOPITT/ASTER product data are pushed to simulated SCF, based on subscriptions

Provide production planning and processing

- All PGE executions in Demo are implemented through Planning (PLS) and Data Processing (DPS), with executions triggered by data arrivals

SCENARIO



Acknowledgments



Special thanks to ASTER, MOPITT, and MISR instrument teams for their support with PGEs and data.